

Inhibition and Antimicrobial Activity of *Lavandula stoechas* Essential Oil on *Escherichia coli*, *Staphylococcus aureus* and *Candida albicans*

Yusuf Sıcak^{1*}, Elif Ayşe Erdogan Eliuz², Eyüp Başaran³⁺, Hatice Ulusoy⁴

¹Department of Plant and Animal Production, Vocational School of Koycegiz, Muğla Sıtkı Koçman University, Muğla, Turkey

²Department of Food Technology, Vocational School of Technical Sciences, Mersin University, Mersin, Turkey

³Department of Chemical and Chemical Processing Technologies, Vocational School of Technical Sciences, Batman University, Batman, Turkey

⁴Department of Forestry, Vocational School of Koycegiz, Muğla Sıtkı Koçman University, Muğla, Turkey

*Corresponding author: yusufsicak@mu.edu.tr

+Speaker: eybasaran@gmail.com

Presentation/Paper Type: Oral / Full Paper

Abstract – In the present study, the volatile compounds of essential oil of *Lavandula stoechas* L. (Lavender) was analysed by gas chromatography-mass spectrometry (GC-MS) using the Nist and Willey libraries. It was determined that the main components of *L. stoechas* were linalyl acetate (36.31%) and linalool (34.18%). Inhibition and antimicrobial activity of volatile oil of the plant against *Escherichia coli* (ATCC 25293), *Staphylococcus aureus* (ATCC25925) and *Candida albicans* were examined by microbroth dilution method. *L. stoechas* EO (Essential Oil) exhibited antimicrobial activity with MIC of between 14.8 and 32.5 µg/mL. The lowest antimicrobial activity were determined against *E. coli* in *L. stoechas* EO (MIC= 32.5 µg/mL), while maximum MIC were noticed as 14.8 µg/mL against *C. albicans*. The most cell inhibition were noted as 91.76% in *C. albicans* at the concentration of 4500 µg/mL of *L. stoechas*. The reductions in living cell number were obtain as 0.54 log in *E. coli*, 1.04 log in *S. aureus* and 1.08 log in *C. albicans* at the same concentration of 4.5 µg/mL. This study showed the strong antimicrobial performance of *L. stoechas* EO in detail. This study showed that *L. stoechas* EO inhibits the growth of microorganisms depending on concentration.

Keywords – *Lavandula stoechas*, antimicrobial action, cell inhibition, reduction of living cell

I. INTRODUCTION

Natural products have therapeutic and antimicrobial characteristics and a large quantity of information has been released [1]. In particular, many studies on the antimicrobial action of essential oils have critical significance. Thus, countless studies against pathogenic bacteria have been recorded to determine antimicrobial operations of vital plant oils. *Lavandula* species are the leading ones of these studies. *Lavandula* genus includes herbaceous plants, annuals, and small shrubs, having aromatic parts. Among cultivated in the world, the most known species believed to have medicinal and aromatherapeutic value are *Lavandula angustifolia*, *Lavandula intermedia*, *Lavandula dentata*, *Lavandula officinalis* and *Lavandula stoechas*. Their essential oil have been used for centuries as a therapeutic agent, especially an antimicrobial agent [2]- [4]. Also culinary herbs and food supplements include most Lavender species based on their biological activities. Many studies has performed several investigations on lavender essential oils, mainly on its antifungal, antioxidant, and anti-inflammatory properties [5]- [8].

Lavandula stoechas L. is a member of Lamiaceae family and distributed in all Mediterranean regions. It is rather important in medical treatment as anticonvulsant, sedative, and antispasmodic activities. In addition, *L. stoechas* is used in perfumery and cosmetics due to consist of desired aromatic components [9], [10]. However, no paper was found reporting the reduction of survival and inhibition percentage

in pathogen microorganisms on different concentration of *L. stoechas* essential oil. Thus, this study focuses on the key issues such as survival of *E. coli*, *S. aureus* and *C. albicans* in natural antimicrobial agent environments and the rate of inhibition of *L. stoechas* EO.

II. MATERIALS AND METHOD

A. Plant material and Essential oil extraction

L. stoechas. were collected from Fethiye region of Muğla, Turkey in 2018. They were identified and confirmed by comparing it with the specimen located at the Herbarium of Biology, Faculty of Science, Muğla Sıtkı Koçman University, Turkey. The HpEO and LsEO extractions were performed from approximately 100 g of the dried *L. stoechas* via hydrodistillation method for 2 hours from several matrix.

B. Chemical composition

The qualitative and quantitative composition of essential oil analysis were conducted at Giresun University central Research Laboratories Application and Research Center by GC-MS 7890A-(5975C inert MSD) instrument equipped with an Agilent 19091S-433 column (30m x 250 µm film x 0.25 µm thickness). Helium was used as a carrier gas. The temperature was raised from 60°C to 225°C by an increase of 3°C/minutes and then 25 minutes of waiting time were implemented during the analysis. Injection port temperature was 250°C, while detector temperature was 260°C. Characterization of *L. stoechas* EO components was

based on the library (Wiley and NIST) comparison with the mass spectra of the injected essential oil samples.

C. Antimicrobial Activity

The antimicrobial activity of essential oil of *L. stoechas* was researched on several pathogens, namely *Escherichia coli*, *Staphylococcus aureus* and *Candida albicans* using and modified spectrophotometric microdilution technique. Firstly, the inoculums of microorganisms were prepared in 4 ml Tryptic Soy Broth for bacteria, 4 ml Sabouraud Dextrose Broth for yeast and incubated at 37°C, overnight. After 24 hours, the culture suspensions were adjusted to 0.5 McFarland Standard Turbidity and stored at +4°C until use [11].

D. Spectrophotometric Microdilution Technique

The experiment were performed on 96-well microtiter plates and firstly 50 µL of Mueller Hinton Broth (MHB) medium were added into all wells. Two-fold serial dilutions (50 µL) of *L. stoechas* (4.65 mg/mL) was made on all x-axis along of elisa plate. Columns 11 and 12 were used as negative and positive controls. Finally, 5 µL culture of microorganisms was inoculated on all wells except medium control wells. All plates were incubated for 24 hours at 37°C, turbidity for bacteria was measured at 600 nm, 415 nm for yeast. The optical density was read before, T0 and T24 after 24-hour incubation for MIC assessment. MIC was calculated for each plate using the following formula: The OD was subtracted from the OD for each replicate at T24 for each replicate. The Percent growth = (ODtest /OD control)x100. Percent Inhibition = 100-(OD test well/OD of corresponding control well)x100 for each row of the 96-well plate. We calculated MIC (the lowest concentration of test material which results in 99.9% inhibition of growth) using the R² formula on inhibition curve [12], [13].

E. Survival under *L. stoechas* Essential Oil

Survival of *Escherichia coli*, *S. aureus* and *C. albicans* were investigated by broth microdilution method in vitro exposure to *L. stoechas* EO. The percentages of cell viability counts were used to assess logarithmic reduction in cell number compared with negative control (100% survivor). To the formule: The reduction of living cell = The cell number in negative control well (MHB + microorganism) - The cell number in corresponding well (MHB + microorganism + corresponding amount of *L. stoechas* EO).

F. Statistical analysis

Survival All data on antimicrobial activity assay studies were the averages of triplicate analysis. Data were recorded as mean ± SEM (standard error of the mean). Significant differences between means were determined by Tukey HSD (SPSS 25; post hoc-one way ANOVA) test and *p* values <0.05 were regarded as significant

III. RESULTS

The components of EO extracted from *L. stoechas* with their retention time (RT) and area (%) were listed in Table 1. In the present study, linalyl acetate (36.31%), linalool (34.18%), camphor (5.38%), 1,8-cineole (3.98%) and borneol (3.11%) were the major component in the essential

oil of *L. stoechas* aerial parts, followed by *lavandulyl acetate* (1.92%), caryophyllene (1.89%), hexyl butyrate (1.45%), α -terpineol (1.31%), β -farnesene (1.23%), geranyl acetate (1.10%) and less amounts than 1% with camphene, 3-octanone, 3-carene, *cis*-ocimene, β -ocimene, neryl acetate, germacrene-D, caryophyllene oxide, α -bisabolol, elaidic acid.

Table 1. Chemical composition of *L. stoechas* essential oil. RT: Retention Time. Quantity (%): more than 0.01.

RT	Compound	%	RT	Compound	%
9.913	Camphene	0.20	24.790	Linalyl acetate	36.31
11.601	3-octanone	0.33	25.878	Lavandulyl acetat	1.92
13.512	3-carene	0.85	28.945	Neryl acetate	0.61
13.747	1,8-cineole	3.98	29.786	Geranyl acetate	1.10
13.947	<i>cis</i> -ocimene	0.70	31.251	Caryophyllene	1.89
14.422	β -ocimene	0.75	32.652	β -farnesene	1.23
17.730	Linalool	34.18	33.682	Germacrene -D	0.31
19.366	Camphor	5.38	38.031	Caryophyllene oxide	0.63
20.333	Borneol	3.11	42.237	α -bisabolol	0.81
21.323	Hexyl butyrate	1.45	57.537	Elaidic	0.33
21.483	(±)- α -terpineol	1.31	Total		99.22

The 24 hour incubation of *L. stoechas* EO with microorganisms was found to be statistically significant in terms of MIC (*p*<0.05) (Table 2). All microorganisms were found to be sensitive to the essential oil. The MIC of *L. stoechas* EO (32.5 µg/mL) against *E. coli* was found to be statistically significant than *L. stoechas* EO (14.8 µg/mL) on *C. albicans*. The moderately antimicrobial activity were determined against *E. coli* and *S. aureus* with MIC= 32.5 µg/mL and MIC= 21.3 µg/mL, respectively, while maximum MIC were noticed as 14.8 µg/mL against *C. albicans* in *L. stoechas*.

Table 2. MIC of *L. stoechas* against tested microbial strains by microdilution method. For positive control: ampicillin (for bacteria) and fluconazole (for yeast) were used as positive control (128 µg/mL). The average MICs were expressed with the standard deviation (±) and significance level (*p*≤0.05) and a: same signs differ statistically at the 0.05 level.

Microorganism/MIC	<i>L. stoechas</i>	Antibiotics
<i>E. coli</i> (-)	32.5 ^a ±3.6	64±3.9
<i>S. aureus</i> (+)	21.3±0.8	8±2.8
<i>C. albicans</i>	14.8 ^a ±1.3	128±4.8

As shown in Table 3, *L. stoechas* EO revealed different inhibition activities towards the three microorganism cells investigated. Inactivation of *E. coli*, *S. aureus* and *C.*


albicans by increased doses of the essential oil was similar in that caused an increase cell death. In general, a dose-dependent decrease in the survival of the microorganisms was observed. The applied essential oil at doses of 8.8, 17.6, 35.2, 70.3, 140.6, 281.3, 562.5, 1125, 2250 and 4500 µg/mL led to inhibition between 3.56% and 71.41% for *E. coli*, 1.17% and 90.90% for *S. aureus*, -5.09% and 91.76% for *C. albicans* ($p < 0.05$). At a concentration of 4.5 mg/mL, the inhibition by *L. stoechas* were 71.41% for *E. coli*, 90.90% for *S. aureus*, 91.76% for *C. albicans*.

Table 3. The logarithmic reduction in cell population (R) and dose-dependent cell inhibition in *E. coli* (E), *S. aureus* (S) and *C. albicans* (C) (%) in the presence of *L. stoechas* essential oil during exposure between the concentration of 4500 µg/mL and 8.789 µg/mL for 24 h.

		Concentration									
		8.78	17.57	35.15	70.31	140.6	281.2	562.5	1125	2250	4500
E	%	3.56	7.19	37.28	56.34	64.86	67.36	68.41	69.09	70.55	71.41
	R	0.02	0.03	0.20	0.36	0.45	0.49	0.50	0.51	0.53	0.54
S	%	1.17	-3.99	-2.26	-2.42	-2.69	-6.08	-4.81	27.71	50.40	90.90
	R	0.01	-0.02	-0.01	-0.01	-0.01	-0.03	-0.02	0.14	0.30	1.04
C	%	-5.09	-3.06	6.83	5.10	5.13	10.48	13.51	28.98	57.98	91.76
	R	-0.02	-0.01	0.03	0.02	0.02	0.05	0.06	0.15	0.38	1.08

The applied *L. stoechas* essential oil at doses of 8.8, 17.6, 35.2, 70.3, 140.6, 281.3, 562.5, 1125, 2250 and 4500 µg/mL led to reduction in live cell count between 0.02 and 0.54 log cfu/mL for *E. coli*, 0.01 and 1.04 cfu/mL for *S. aureus*, -0.02 and 1.08 cfu/mL for *C. albicans* ($p < 0.05$). The high reduction of living cell were obtain as 1.08 log in *C. albicans* at the concentration of 4.5 mg/mL of *L. stoechas*. The others also: It was 1.04 log in *S. aureus* and 0.54 log in *E. coli* at the same concentration. *E. coli* cell population during exposure in the essential oil (2.25 mg/mL) poured medium decreased 0.53 log according to the control experiments which is without addition of essential oil. *S. aureus* and *C. albicans* cell population decreased 0.30 log and 0.38 log, respectively. As for inactivation of *E. coli*, *S. aureus* and *C. albicans* under the same doses of *L. stoechas* EO, the 1125 mg/mL EO decreased the initial count of 0.51, 0.14, 0.15 log cfu/mL in the control samples (2 log cfu/mL), respectively ($p < 0.05$). It was 0.50, -0.02, 0.06 log cfu/mL for *E. coli*, *S. aureus* and *C. albicans* in the 562.5 mg/mL EO, respectively (Figure 1).

Figure 1. The logarithmic reduction in cell population (line) and dose-dependent cell inhibition in *E. coli*, *S. aureus* and *C. albicans* in the presence of *L. stoechas* essential oil during exposure between the concentration of 4500 µg/mL and 8.789 µg/mL for 24 h. The cell number in control samples: 2 log cfu/mL



IV. DISCUSSION

Although the most widely common chemotype of *L. stoechas* is the camphor-fenchone, we noted the linalyl acetate, linalool, camphor which have strong pharmacological activities such as antimicrobial, anti-inflammatory [14]. Many authors have attributed this chemical difference to differences in habitat, season and plant origin [15], [16]. In existing literature, the oil of *L. stoechas* were noted mostly consists of linalyl acetate, 1,8-cineole camphene and fenchone [17]. Soyulu et al. (2006) obtained that the main components of *L. stoechas* were 1,8-Cineole (35.5%), sabinene (15.0%), α -terpineyl acetate (14.2%), α -pinene (7.5%) [18]. In another study, the major compounds in *L. stoechas* EO were reported to be fenchone (11.27-37.48%), camphor (1.94-21.8%), 1,8-cineole (0.16-8.71%), and viridiflorol (2.89-7.38%) [19]. The GC-MS profile depicted that fenchone (55.79%), camphor (18.18%), 1,8-cineole (8.03%), and myrtenyl acetate (6.25%) were the major components in *L. stoechas* EO [20]. Similarly, the featured constituents of the essential oils from various populations of *L. stoechas* L. ssp. *stoechas* in Greece were α -pinene, 1,8-cineole, fenchone, camphor and myrtenyl acetate, all components of different amounts due to different habitat [21]. The component of fenchone from the stems/leaves and flowers of *L. stoechas* L. ssp. *stoechas* mostly was extracted on varying proportions such as 13.13%, 27.08%, 52.60% [16].

Previous studies showed the essential oils of Lavender species is widely effective against the growth of a wide range of pathogens, particularly, *E. coli*, *S. aureus* and *C. albicans* [22], [23]. These effect comes from the components such as linalyl acetate, linalool, camphor, fenchone, caryophyllene, hexyl butyrate, α -terpineol, β -farnesene that it possesses. The MIC values of *L. angustifolia* essential oil extracted by hydrodistillation and having high linalool ratio (52.59%) were found to be over 1.25 mg/mL against *E. coli*, *S. aureus* and *C. albicans* [24]. In another study, the essential oils of *L. angustifolia* and *Lavandula intermedia* which include great amount of caryophyllene and camphor, respectively, effectively inhibited against *Staphylococcus aureus*, *E. coli* [4]. Also, Soyulu et al. (2006) showed that Lavender EO were strong fungicidal at relatively above concentrations of 12.8 µg/mL [18].

In our study, linalyl acetate (36.31%), linalool (34.18%), camphor (5.38%) were be found high level and they were effective to all pathogens. Similarly, Arıdoğan et al (2002) demonstrated that *L. hybrida* contain linalool (32.8%), linalylacetate (29.9%) and had anti-staphylococcus activity (8 mm) and had no anti- *E. coli* activity (0 mm) by disc diffusion method [25]. In another study, the *L. stoechas* EO exhibited good antimicrobial activity against *E. coli*, *S. aureus* and *C. albicans* at the concentrations (MICs) ranging from 0.16 to 11.90 mg/mL [19]. In addition, it was reported that linalyl acetate showed low antibacterial activity with MIC at 7.0-10.0 µg/mL in the microdilution method, while linalool showed bacteriostatic activity at 4.0-7.0 µg/mL and bactericidal effect at 4.0-9.0 µg/mL [26].

L. stoechas EO showed above 70% inhibitory effect at concentration of 4.5 mg/mL on the pathogens. They were 91.76 for *C. albicans*, 90.90% for *S. aureus* and 71.41% for *E. coli*. Normally, the inhibition activity was decreased in the lower concentration of *L. stoechas* EO. The inhibition ratio were 57.98% for *C. albicans*, 50.40% for *S. aureus* and 70.55% for *E. coli* in 2250 mg/mL *L. stoechas* EO. At the concentration of 1125 mg/mL *L. stoechas* EO, it was 28.98% for *C. albicans*, 27.71% for *S. aureus* and 69.09% for *E. coli*. Remarkably, inhibition activity of *L. stoechas* EO (562.5 mg/mL) were not reported on *S. aureus*. However, there were significantly activity were noted 68.41% for *E. coli* and 13.51% for *C. albicans* at the same concentration. Few studies have reported the inhibition rate on pathogens of essential oil of *L. stoechas*. *Staphylococcus aureus* was reported to be rather sensitive to *Lavandula stoechas* EO which its main component were fenchone (68.2%) and camphor (11.2%) [17]. Linalool and linalyl acetat caused more than 50% inhibition on *C. albicans* [27].

As can be seen, the logarithmic reduction in the number of cells with percent inhibition is shown in the same graph. The count of viable cells decreased as the concentration of the essential oil increased. For inactivation of microorganisms, the degree of inhibitory effects of *L. stoechas* EO was *E. coli*>*C. albicans*>*S. aureus* at mostly high concentration, for example, 4.5 mg/mL ($p<0.05$). There is a study about this. Dadalıoğlu and Evrendilek (2004) reported that Spanish *L. stoechas* essential oil with the concentration of 5 and 80 µL/mL reduced the live cell count around 4 log cfu/mL. As for inactivation of *S. aureus* under the 5 and 10 µL/mL doses of the lavender decreased the initial count of about 3 log cfu/mL [20].

V. CONCLUSION

In conclusion, the inhibition and antimicrobial potential of *L. stoechas* against *E. coli*, *S. aureus* and *C. albicans* were demonstrated. All microorganisms were sensitive to the oil and the inhibition ratio and also the reduction in cell count were reported for different concentrations of the oil. The results in our study will shed light on future research.

ACKNOWLEDGMENT

Thanks to Giresun University Central Research Laboratories Application and Research Center for GC-MS.

REFERENCES

- [1] A. Lopez, J.B. Hudson, G.H.N. Towers, "Antiviral and antimicrobial activities of Colombian medicinal plants". *J Ethnopharmacol.*, vol. 77, pp. 189-96, Oct. 2001.
- [2] N.Y.O. Muyima, G. Zulu, T. Bhengu, D. Popplewell, "The potential application of some novel essential oils as natural cosmetic preservatives in an aqueous cream formulation," *Flavour Fragr. J.*, vol. 17, pp. 258-266, March. 2002.
- [3] S. De Rapper, G. Kamatou, A. Viljoen, S. Van Vuuren, "The in vitro antimicrobial activity of *Lavandula angustifolia* essential oil in combination with other aroma-therapeutic oils," *Evid-Based Compl Altern Med.*, Vol. 2013:852049, Apr. 2013.
- [4] C. Jianu, G. Pop, A.T. Gruiu, F.G. Horhat, "Chemical composition and antimicrobial activity of essential oils of lavender (*Lavandula angustifolia*) and lavandin (*Lavandula x intermedia*) grown in Western Romania," *Int J Agric Biol*, Vol. 15, pp. 772-776, March. 2013.
- [5] V. Hajhashemi, A. Ghannadi, B. Sharif, "Anti-inflammatory and analgesic properties of the leaf extracts and essential oil of *Lavandula angustifolia* Mill.," *J Ethnopharmacol.*, vol. 89, pp. 67-71, Nov. 2003.
- [6] I. Spiridon, S. Colceru, N. Anghel, C.A. Teaca, R. Bodirlau, A. Armatu, "Antioxidant capacity and total phenolic contents of oregano (*Origanum vulgare*), lavender (*Lavandula angustifolia*) and lemon balm (*Melissa officinalis*) from Romania," *Nat Prod Res*. Vol. 25, pp. 1657-61, Jun. 2011.
- [7] M. Zuzarte, M.J. Gonçalves, C. Cavaleiro, J. Canhoto, L. Vale-Silva, M.J. Silva, E. Pinto, L. Salgueiro, "Chemical composition and antifungal activity of the essential oils of *Lavandula viridis* l'Her.," *J Med Microbiol.*, Vol. 60, pp. 612-618, May. 2011.
- [8] M. Zuzarte, L. Vale-Silva, M.J. Gonçalves, C. Cavaleiro, S. Vaz, J. Canhoto, E. Pinto, L. Salgueiro, "Antifungal activity of phenolic-rich *Lavandula multifida* L. Essential oil," *Eur J Clin Microbiol Infect Dis.*, Vol. 31, pp. 1359-66, Oct. 2012.
- [9] K. Canli, A. Yetgin, A. Benek, M.E. Bozyel, E.M. Altuner, "In Vitro Antimicrobial Activity Screening of Ethanol Extract of *Lavandula stoechas* and Investigation of Its Biochemical Composition," *Adv Pharmacol Sci*, Volume 2019, ID 3201458, janu. 2019.
- [10] A.H. Gilani, N. Aziz, M.A. Khan, F. Shaheen, Q. Jabeen, B.S. Siddiqui, J.W. Herzig, "Ethnopharmacological evaluation of the anticonvulsant, sedative and antispasmodic activities of *Lavandula stoechas* L.," *J Ethnopharmacol.*, vol 71, pp. 161-167, Nov. 2000.
- [11] *Dalynn Biologicals*, McFarland Standard, Cat no: TM50-TM60, 2014.
- [12] T. Patton, J. Barrett, J. Brennan, N. Moran, "Use of a spectrophotometric bioassay for determination of microbial sensitivity to manuka honey," *J Microbiol Methods.*, Vol. 64, pp. 6484-6495, Apr. 2006.
- [13] Y. Sıcak, E.A. Erdogan Eliuz, "Determination of the phytochemical profile, in vitro the antioxidant and antimicrobial activities of essential oil from *Arbutus andrachne* L. wood growing in Turkey," *Turkish J Forestry.*, Vol. 20, pp. 57-61, March. 2019.
- [14] A.T. Peana, P.S. D'Aquila, F. Panin, G. Serra, P. Pippia, M.D.L. Moretti, "Anti-inflammatory activity of linalool and linalyl acetate constituents of essential oils.," *Phytomedicine.*, Vol., 9, pp. 721-726. 2002.
- [15] A.C. Gören, G. Topçu, G. Bilsel, M. Bilsel, Z. Aydoğmuş, J.M. Pezzuto, "The chemical constituents and biological activity of essential oil of *Lavandula stoechas* ssp. *stoechas*," *Z Naturforschung C.*, Vol., 57, pp. 9-10, June. 2002.
- [16] A. Angioni, A. Barra, V. Coroneo, S. Dessi, P. Cabras, "Chemical composition, seasonal variability, and antifungal activity of *Lavandula stoechas* L. ssp. *stoechas* essential oils from stem/leaves and flowers," *J Agric Food Chem.*, Vol. 54, pp. 4364-70, May. 2006.
- [17] N. Bouzouita, F. Kachouri, M. Hamdi, M.M. Chaabouni, R. Ben Aissa, S. Zgoulli, "Volatile constituents and antimicrobial activity of *Lavandula stoechas* L. oil from Tunisia," *J Essent Oil Res.*, Vol. 17, pp. 584-586., Nov. 2005.
- [18] E.M. Soyulu, S. Soyulu, S. Kurt, "Antimicrobial activities of the essential oils of various plants against tomato late blight disease agent *Phytophthora infestans*," *Mycopathologia.*, Vol. 161, pp. 119-128, Oct. 2006.
- [19] T. Benabdelkader, A. Zitouni, Y. Guitton, F. Jullien, D. Maitre, H. Casabianca, L. Legendre, A. Kameli, "Essential oils from wild

- populations of algerian *Lavandula stoechas* L. Composition, chemical variability, and in vitro biological properties," *Chem Biodivers.*, Vol. 8, pp. 937-953, May, 2011
- [20] I. Dadalıođlu, G.A. Evrendilek, "Chemical compositions and antibacterial effects of essential oils of Turkish oregano (*Origanum minutiflorum*), bay laurel (*Laurus nobilis*), Spanish lavender (*Lavandula stoechas* L.), and fennel (*Foeniculum vulgare*) on common foodborne pathogens," *J Agric Food Chem.*, Vol., 52(26): 8255-60, Nov. 2004.
- [21] M. Skoula, C. Abidi, E. Kokkalou, "Essential oil variation of *Lavandula stoechas* L. ssp. *stoechas* growing wild in Crete (Greece)," *Biochem Syst Ecol.*, Vol. 24, pp. 255-260, Apr. 1996.
- [22] K.A. Hammer, C.F. Carson, T.V. Riley, "Antimicrobial activity of essential oils and other plant extracts," *J Appl Microbiol.*, Vol. 86, pp. 985-990. Dec. 1999.
- [23] L. Hui, L. Huan, L. XiaoLan, Z. AiGuo, "Chemical composition of lavender essential oil and its antioxidant activity and inhibition against rhinitis-related bacteria," *Afr J Microbiol Res.*, Vol. 4, pp. 309-313. Jan. 2010.
- [24] L.T. Danh, L.N. Han, N.D.A. Triet, J. Zhao, R. Mammucari, N. Foster, "Comparison of Chemical Composition, Antioxidant and Antimicrobial Activity of Lavender (*Lavandula angustifolia* L.) Essential Oils Extracted by Supercritical CO₂, Hexane and Hydrodistillation," *Food Bioprocess Tech.*, Vol. 6, pp. 3481-3489, Dec. 2013.
- [25] B.C. Aridođan, H. Baydar, S. Kaya, M. Demirci, D. Özbařar, E. Mumcu, "Antimicrobial activity and chemical composition of some essential oils," *Arch Pharm Res.*, Vol. 25, pp. 860-864, Dec. 2002.
- [26] M. Sokovic, J. Glamođlija, P.D. Marin, D. Brkić, L.J.L.D. Van Griensven, "Antibacterial effects of the essential oils of commonly consumed medicinal herbs using an in vitro model," *Molecules*, Vol. 15, pp. 7532-7546, Oct. 2010.
- [27] G.B. Zore, A.D. Thakre, S. Jadhav, S.M. Karuppayil, "Terpenoids inhibit *Candida albicans* growth by affecting membrane integrity and arrest of cell cycle," *Phytomedicine*, Vol. 18, pp. 1181-1190. Oct. 2011.